

MDU M.Sc Entrance: Model Test

Chemistry

❖ Question Paper

All questions are compulsory (One mark each)

Total Marks: 100 (1.5 Hours)

Q.1 $[\text{CoCl}_4]^{2-}$ shows a deep blue colour because of

- (a) Metal to ligand charge transfer transition.
- (b) Ligand to metal charge transfer transition.
- (c) Spin allowed and Laporte forbidden d-d transition.
- (d) Spin allowed and Laporte allowed d-d transition.

Q.2 The violet colour of iodine vapour is due to

- (a) $n \rightarrow n^*$ transition
- (c) $\pi \rightarrow \pi^*$ transition
- (c) $n \rightarrow \sigma^*$ transition
- (d) $\sigma \rightarrow \pi^*$ transition

Q.3 Choose the correct statement among the following

- (a) Diamond has lower thermal and electrical conductivities compared to graphite.
- (b) Diamond has similar thermal and electrical conductivities compared to graphite.
- (c) Diamond has higher thermal conductivity but lower electrical conductivity compared to graphite
- (d) Diamond has the same thermal but lower electrical conductivity compared to graphite.

Q.4 Which of the following is a nido-borane?

- (a) B_4H_{10}
- (b) B_5H_9
- (c) $[\text{B}_6\text{H}_6]^{2-}$
- (d) B_5H_{11}

Q.5 Among the three types of orbital s, p, d, and f,

- (a) Both p and f orbitals have centre of symmetry
- (b) Both p and d orbitals have centre of symmetry

- (c) Only d orbitals have centre of symmetry (d) f orbitals alone have centre of symmetry

Q.6 The absorbance of solution having 20% transmittance is

- (a) 0.301 (b) 0.699 (c) 1.301 (d) 1.699

Q.7 The active site of enzyme nitrogenase contains

- (a) Mo (b) Mn (c) Fe (d) Cu

Q.8 Which one of the following is a free radical?

- (a) CO (b) CN^- (c) NO (d) CS

Q.9 Choose the $16 e^-$ complex from the following:

- (a) $\text{Ni}(\text{CO})_4$ (b) $\text{Rh}(\text{PPh}_3)_3\text{Cl}$ (c) $\text{Fe}(\text{CO})_5$ (d) $(\eta^6\text{-C}_6\text{H}_6)_2\text{Cr}$

Q.10 The species having metal-metal bond is:

- (a) $\text{Mn}_2(\text{CO})_{10}$ (b) $\text{Al}_2(\text{CH}_3)_6$ (c) $\text{V}_2(\text{CO})_{12}$ (d) $\text{Al}_2(\text{OPr})_{12}$

Q.11 The only molecule having bridging oxygen is

- (a) Phosphorus trioxide (b) Phosphorus pentoxide
(c) Cyclic tetraphosphate (d) Pyrophosphate

Q.12 The coordination number of phosphorus in $[\text{PMo}_{12}\text{O}_{40}]^{3-}$ is

- (a) 2 (b) 4 (c) 5 (d) 6

Q.13 Using phenolphthalein as the indicator, which of the following titration is possible:

- (a) Acetic acid with pyridine (b) Oxalic acid with sodium hydroxide
(c) Hydrochloric acid with aniline (d) Sulphuric acid with aqueous ammonia

Q.14 Which of the following species is ESR-active?

- (a) VOSO_4 (b) $\text{K}_2\text{Cr}_2\text{O}_7$ (c) KMnO_4 (d) $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$

Q.15 Large deviation from Trouton's rule is observed for systems which are

- (a) Having more ordered structure (b) Having more disordered structure
(c) Having low melting points (d) Having low boiling points

Q.16 The concentration of a reactant decreases linearly with time. What is the order of the reaction?

- (a) 1st order (b) Fractional order (c) 2nd order (d) Zero order

Q.17 I for C^{-13} is:

- (a) 1 (b) $1/2$ (c) $3/2$ (d) 2

Q.18 The number of rotational degrees of freedom of CO_2 is

- (a) One (b) Two (c) Three (d) Four

Q.19 The magnitude of the nuclear spin angular momentum of a nuclei is $\sqrt{15/2}\hbar$ units. The value of I is

- (a) $5/2$ (b) $1/2$ (c) 1 (d) $3/2$

Q.20 Which of the following transitions in the electronic spectrum of a homonuclear diatomic molecule is forbidden

- (a) $\Sigma_u^+ \rightarrow \Sigma_g^+$ (b) $\Sigma_g^+ \rightarrow \Pi_u^+$ (c) $\Sigma_u^+ \rightarrow \Pi_g^+$ (d) $\Sigma_g^+ \rightarrow \Delta_u$

Q.21 The diffraction pattern of a cubic solid has an intense 110 Bragg reflection, but the 100 and 111 Bragg reflections are absent. The structure of the solid is

- (a) Body-centered cubic (b) Primitive cubic
(c) Face-centered cubic (d) Edge-centered cubic

Q.22 The logarithmic conductivity of a crystalline solid shows a linear variation with inverse temperature ($1/T$). The band gap may be obtained from

- (a) Slope of the plot. (b) Intercept on the conductivity axis.
(c) Intercept on the temperature axis. (d) Inverse slope.

Q.23 C = C frequency in oct-4-ene appears at:

- (a) $1680-1600\text{ cm}^{-1}$ (very weak) (b) $1680-1600\text{ cm}^{-1}$ (strong)
(c) $1680-1600\text{ cm}^{-1}$ (m) (d) No peak in this region of $1680-1600\text{ cm}^{-1}$

Q.24 The spatial part of hydrogen molecular wave function in the simplest molecular orbital theory is given by σ_g^2 where σ_g is a normalized linear combination of two hydrogen 1s orbitals. Which of the following is true about the above wave function?

- (a) It contains only covalent terms. (b) It includes only a small amount of ionic terms.
(c) It contains only ionic terms. (d) It over estimates the ionic terms.

Q.25 A $2p_z$ orbital of hydrogen atom is an eigenfunction of

- (a) H only (b) H and L^2 only (c) H, L^2 and L_z only (d) H, L^2 , L_z and L_x


Q.26 By a reversible process, we mean one that always

- (a) Takes infinite time for completion (b) Satisfies $\Delta S(\text{universe}) = 0$
(c) Satisfies $\Delta G = 0$ (d) Gives the minimum work

Q.27 A hydrogenic 3p orbital has the following form of the radial wavefunction ($\alpha_1 = \text{constant}$):

- (a) $r(\alpha_1 - r)e^{-\alpha_2 r}$ (b) $re^{-\alpha_3 r}$
 (c) $r(\alpha_4 - r)(\alpha_5 - r)e^{-\alpha_6 r}$ (d) $r^3 e^{-\alpha_3 r}$

Q.28 IUPAC name for the compound given below is



- (a) E-5-ethylhept-5-en-2-one (b) Z-5-ethylhept-5-en-2-one
 (c) E-3-ethylhept-2-en-6-one (d) Z-3-ethylhept-2-en-6-one

Q.29 The most suitable reagent for the following transformation is



- (a) NaBH_4 (b) B_2H_6 (c) Zn-Hg/HCl (d) $\text{NH}_2\text{NH}_2/\text{HCl}$

Q.30 The major product formed in the reaction of 2-methyl but-3-en-2-ol with HBr is

- (a) (b) (c) (d)

Q.31 Among dimethylcyclobutanes, which one can exhibit optical activity?

- (a) cis-1,2-dimethylcyclobutane (b) trans-1,2-dimethylcyclobutane
 (c) cis-1,3-dimethylcyclobutane (d) trans-1,3-dimethylcyclobutane


Q.32 The monomer of biopolymer DNA is a

- (a) Nucleotide (b) Amino acid (c) Disaccharide (d) Fatty acid


Q.33 The order of chemical shifts (δ value) in the ^1H NMR spectrum of crotonaldehyde is

- (a) Olefinic > CHO > Me (b) CHO > Me > olefinic (c) CHO > Olefinic > Me (d) Olefinic > Me > CHO

Q.34 The product formed in the reaction given below is




Q.35 The major product formed in the reaction given below is



Q.36 The conversion of excited singlet state (S_1) of a molecule to triplet state (T_1) is known as


- (a) Fluorescence (b) Phosphorescence (c) Intersystem crossing (d) Internal conversion

Q.37 The decreasing order of stability of the free radicals A, B and C is




- (a) $A > B > C$ (b) $C > A > B$ (c) $B > A > C$ (d) $A > C > B$

Q.38 The major product formed in the reaction given below:




Q.39 The rates of keto-enol tautomerism in the ketones A-C, given below are in the order



- (a) $A > B > C$ (b) $A > C > B$ (c) $C > A > B$ (d) $C > B > A$

Q.40 The reaction given below is an example of



- (a) Aldol condensation (b) Knoevenagel condensation
(c) Dieckmann condensation (d) Acyloin condensation

Q.41 The covalent radii vary gradually in the Periodic Table. From the orders given below for such radii, the correct ones are

(a) $Ce > Lu$, (b) $Co > Ti$, (c) $Sr > Ca$, (d) $I > Se$

(a) (a) and (b) only (b) (a) and (c) only (c) (a), (c) and (d) only (d) (b), (c) and (d) only

Q.42 The pair of gaseous molecules/ions having tetrahedral structure is

(a) $SnCl_4$, PH_4^+ (b) $SnCl_4$, XeF_4 (c) ICl_4^- , PH_4^+ (d) $SnCl_4$, ICl_4^-

Q.43 Consider the following:

Volumetric method for Ag(I)	Indicator used
(a). Fajan method	Chromate
(b). Mohr's method	Fluorescein
(c). Vohlard method	ferrie salt

The method and indicator matches correctly in

(a) (a) and (b) only (b) 2. (b) and (c) only (c) (c) only (d) (b) only

Q.44 An unknown lead solution has diffusion current of $1.0 \mu A$. To a 10 ml of this solution 0.5 ml of 0.04 M lead solution is added. The diffusion current of the spiked solution is $1.50 \mu A$. The concentration of the unknown lead solution is

(a) 0.0020 M (b) 0.0050 M (c) 0.0035 M (d) 0.0010 M

Q.45 The ^{32}P radio isotope, used in leukemia therapy, has $t_{1/2} = 14.26$ days. What % of ^{32}P remains after 35 days?

(a) 30% (b) 8% (c) 81.7% (d) 18.3%

Q.46 On a 30 cm column, the t_R of A and B respectively are 16.40 and 17.63 minutes. The t_0 of the column is 1.30 minutes. The peak width at base lines for A and B are 1.11 and 1.21 minutes respectively. The column resolution R_s is

(a) 1.06 (b) 1.23 (c) 2.12 (d) 2.23

Q.47 Which one of the following pairs of electronic configurations of high-spin transition metal ions (3d) in an octahedral field undergoes a substantial John-Teller distortion:

- (a) d^3, d^9 (b) d^4, d^9 (c) d^5, d^9 (d) d^6, d^9

Q.48 Which one of the following pairs consists of a good oxidizing and a good reducing agent respectively:

- (a) Ce(IV), Ln(III) (b) Ln(III), Eu(II) (c) Ce(IV), Eu(II) (d) Ln(III), Ce(III)

Q.49 Which one of the pairs of following statements about reduction of $[\text{CoCl}(\text{NH}_3)_5]^{2+}$ By Cr(II) is correct:

- (A). Reactant $[\text{CoCl}(\text{NH}_3)_5]^{2+}$ has non-labile coordination sphere
(B). Reaction proceeds by outer-sphere mechanism
(C). Reactant $[\text{CoCl}(\text{NH}_3)_5]^{2+}$ has labile coordination sphere
(D). Reaction proceeds by inner-sphere mechanism

- (a) (A) and (B) (b) (A) and (D) (c) (C) and (B) (d) (C) and (D)

Q.50 Hemocyanin contains

- (a) A dinuclear copper core and binds dioxygen in the cuprous state.
(b) A dinuclear copper core and binds dioxygen in the cupric state.
(c) A mononuclear copper core and binds dioxygen in the cuprous state.
(d) A mononuclear copper core and binds dioxygen in the cupric state.

Q.51 The ^{31}P NMR spectrum of $\text{PF}_4\text{N}(\text{CH}_3)_2$ at room temperature and low temperature (173K) respectively shows (assume that N and H do not couple):

- (a) Triplet and quintet (b) Quintet and triplet
(c) Quintet and triplet of triplets (d) Triplet and triplet of triplets

Q.52 The number of hyperfine lines in the EPR spectrum of a one electron reduced product of $[\text{Co}_3(\text{CO})_9\text{Se}]$ ($I=7/2$ for Co nucleus) is:

- (a) 8 (b) 15 (c) 22 (d) 1

Q.53 The highest oxidation state of a metal in the following compounds is :

$(\eta^6\text{-C}_6\text{H}_6)_2\text{Cr}$, $\text{Mn}(\text{CO})_5\text{Cl}$, $\text{Na}_2[\text{Fe}(\text{CO})_4]$, $\text{K}[\text{Mn}(\text{CO})_5]$ and $\text{K}[\text{Mo}(\text{CO})_5\text{Br}]$

- (a) 1 (b) 2 (c) -1 (d) -2

Q.54 The maximum number of valence electrons of a metal in these complexes are: $\text{Mn}_2(\text{CO})_{10}$, $(\eta^5\text{-C}_5\text{H}_5)\text{Mo}(\text{CO})_3\text{Cl}$, $(\eta^5\text{-C}_5\text{H}_5)_2\text{Ni}$, and $(\eta^5\text{-C}_5\text{H}_5)_2\text{TiCl}_2$

- (a) 16 (b) 18 (c) 20 (d) 22

Q.55 Olefin hydrogenation using Wilkinson's catalyst initiates with:

- (a) Olefin addition to $\text{Rh}(\text{PPh}_3)_2\text{Cl}$.
(b) Olefin addition to $\text{Rh}(\text{PPh}_3)_3\text{Cl}$.
(c) A phosphine dissociation from $\text{Rh}(\text{PPh}_3)_3\text{Cl}$.
(d) A phosphine addition to $\text{Rh}(\text{PPh}_3)_2\text{Cl}$.

Q.56 Although Fe(III) is a better Lewis acid compared to Zn(II), most hydrolytic Enzymes contain Zn(II) at the active site because

- (a) Fe(III) is a redox active ion.
(b) Fe(III) has less abundance compared to Zn(II).
(c) Fe(III) generally makes octahedral complexes while Zn(II) makes tetrahedral complexes.
(d) Zn(II) makes kinetically labile complexes.

Q.57 Considering the two complexes (A) $[\text{Ni}(\text{H}_2\text{O})_6]^{2+}$ and (B) $[\text{Ni}(\text{NH}_3)_6]^{2+}$, the right statement is

- (a) Complex (A) is diamagnetic and complex (B) is paramagnetic.
(b) Complex (A) is paramagnetic and complex (B) is diamagnetic.
(c) Both are paramagnetic.

- (d) Both are diamagnetic.

Q.58 Unlike d-d transitions, the f-f transitions

- (a) Do not change much with change in ligand.
(b) Change significantly with change in ligand.
(c) Appear at low energies i.e., at the near-IR region.
(d) Appear as broad bands.

Q.59 Strongest super acid among the following is a

- (a) Solution of HNO_3 in H_2SO_4 (b) Solution of HClO_4 in H_2SO_4
(c) Solution of SbF_5 in HF (d) Solution of SbCl_5 in HCl

Q.60 Consider the following statements regarding borazine:

- A. It is isoelectronic with benzene.
B. Each nitrogen receives more σ -electron density from neighboring boron than it gives away as a π -donor.
C. It does not undergo addition reactions.
D. Nitrogen retains its basicity and boron its acidity.


The true statements among the above are

- (a) A, C and D (b) A, B and D (c) A and C only (d) B, C, and D

Q.61 For a diffusion-controlled bimolecular reaction, the rate constant (k_D) is proportional to (T = temperature; η = coefficient of viscosity of medium).

- (a) ηT (b) $1/\eta T$ (c) $\frac{T}{\sqrt{\eta}}$ (d) T/η

Q.62 Consider the unimolecular reaction



For which the following remarks were made.

- A. The reaction is second order at low pressure and becomes first order at high pressure.
 B. The reaction is first order at low pressure and becomes second order at high pressure.
 C. The reaction is zero order.

Which of these is/are correct?

- (a) A and B (b) B and C (c) Only C (d) Only A

Q.63 A random distribution of errors obeys the Gaussian form $\sqrt{A/\pi} \exp[-Ax^2]$. The mean and standard deviation of this distribution obeys

- (a) $\langle x \rangle = 0$ and $\sigma_x = \sqrt{2A}$ (b) $\langle x \rangle \neq 0$ and $\sigma_x = 1/\sqrt{2A}$
 (c) $\langle x \rangle = 0$ and $\sigma_x = \sqrt{A}$ (d) $\langle x \rangle = 0$ and $\sigma_x = A$

Q.64 The function $\sin^{-1}x$ is not an acceptable wave function because

- (a) It is not differentiable (b) Its first derivative is not continuous
 (c) It does not cover the entire space (d) It is not a single-valued function

Q.65 The first-order correction to energy for the ground state of a particle-in-a-box due to a perturbation λx would be

- (a) $\lambda L/2$ (b) λL (c) $2\lambda L$ (d) 2

Q.66 The wave function for a particle in one dimensional box is expressed as

- (a) $\frac{\sqrt{2}}{a} \sin \frac{n\pi x}{a}$ (b) $\sqrt{\frac{2}{a}} \sin \frac{n\pi x}{a}$ (c) $\sqrt{\frac{2}{a}} \sin \frac{\pi x}{a}$ (d) $\sqrt{\frac{2}{a}} \sin \frac{2\pi x}{a}$

Q.67 The Boyle temperature is that at which the second virial coefficient of real gas is

- (a) Zero (b) One (c) Four (d) One and half

Q.68 The energy levels of cyclopropene are $\alpha + 2\beta$, $\alpha - \beta$, and $\alpha - \beta$. The delocalization energy in C_3H_3 is

- (a) 2β (b) 0 (c) β (d) 3β

Q.69 The rotational constant (B) of $H^{35}Cl$, $H^{37}Cl$ and $D^{35}Cl$ follow the order

- (a) $H^{35}Cl > D^{35}Cl > H^{37}Cl$ (b) $H^{35}Cl > H^{37}Cl > D^{35}Cl$
 (c) $D^{35}Cl > H^{35}Cl > H^{37}Cl$ (d) $H^{37}Cl > H^{35}Cl > D^{35}Cl$

Q.70 In a crystal, atom A is at the corners of the unit cell, B is at the centre of the cell and the oxygen atoms are at the face-centred positions. What is the formula per unit cell?

- (a) A_8BO_6 (b) ABO_6 (c) A_8BO_3 (d) ABO_3

Q.71 On mixing 100 mL of 0.1 M CH_3COOH and 50 mL of 0.1 M $NaOH$, the pH of the solution will be

- (a) $pK_a + 0.301$ (b) pK_a (c) $pK_a - 0.301$ (d) $pK_a + 0.477$


Q.72 Using the fundamental equation $dA = -SdT - PdV$, the Maxwell relation is

- (a) $\left(\frac{\partial S}{\partial P}\right)_T = \left(\frac{\partial V}{\partial S}\right)_P$ (b) $\left(\frac{\partial S}{\partial V}\right)_T = \left(\frac{\partial P}{\partial T}\right)_V$ (c) $\left(\frac{\partial T}{\partial V}\right)_T = \left(\frac{\partial P}{\partial S}\right)_T$ (d) $\left(\frac{\partial S}{\partial V}\right)_T = \left(\frac{\partial P}{\partial T}\right)_V$

Q.73 The relationship between mean ionic activity coefficient for $Ca_3(PO_4)_2$ and its ions is given by

- (a) $\gamma_{\pm} = \gamma_+^3 \gamma_-^2$ (b) $\gamma_{\pm} = \gamma_+^2 \gamma_-^3$ (c) $\gamma_{\pm}^5 = \gamma_+^3 \gamma_-^2$ (d) $\gamma_{\pm}^5 = \gamma_+^2 \gamma_-^3$

Q.74 Assuming that C-H and C-X bond lengths in



are nearly equal, the molar residual entropy at 0 K is

- (a) 0 (b) $R \ln 2$ (c) $R \ln 3$ (d) $R \ln 6$

Q.75 The contributions to the molar entropy by translational (tr), rotational (rot), vibrational (vib) and electronic (ele) degrees of freedom is in order

- (a) $\text{tr} > \text{rot} > \text{vib} > \text{ele}$ (b) $\text{rot} > \text{vib} > \text{tr} > \text{ele}$ (c) $\text{ele} > \text{vib} > \text{rot} > \text{tr}$ (d) $\text{vib} > \text{rot} > \text{tr} > \text{ele}$

Q.76 A binary mixture of A_2 and B_2 will show negative deviation from Raoult's law when

- (a) A-A and B-B interactions are stronger than A-B.
(b) A-A and B-B interactions are weaker than A-B.
(c) Both A-A and B-B interactions are equal to A-B.
(d) Either A-A or B-B interactions is equal to A-B.

Q.77 In the presence of external magnetic field the transition $^3D_1 \rightarrow ^3P_1$ splits into

- (a) 3 (b) 5 (c) 7 (d) 9

Q.78 Ionic equivalent conductance value for Ca^{2+} is $0.0119 (\text{S m}^2\text{mol}^{-1})$ and for Cl^- is $0.0076 (\text{S m}^2\text{mol}^{-1})$. The correct expected molar conductivity at infinite dilution for CaCl_2 is

- (a) $0.0195 \text{ S m}^2\text{mol}^{-1}$ (b) $0.0271 \text{ S m}^2\text{mol}^{-1}$ (c) $0.0542 \text{ S m}^2\text{mol}^{-1}$ (d) $0.01355 \text{ S m}^2\text{mol}^{-1}$

Q.79 The term symbol for the ground state configuration of NO is

- (a) $^2\Pi_u$ (b) $^2\Pi_g$ (c) $^1\Pi_u$ (d) $^1\Pi_g$


Q.80 The kinetic chain length (ν) is a measure of chain propagation. If the rates of consumption are denoted by R_c and rates of production by R_p ; M and M^* denote the monomer and the active center, respectively. The correct definition of ν is

- (a) $R_c(M)/R_p(M^*)$ (b) $R_p(M^*)/R_c(M)$ (c) $R_c(M^*)/R_p(M)$ (d) $R_c(M)/R_c(M^*)$


Q.81 4-tert-Butylcyclohexanone reduction gives two isomeric alcohols which are

- (a) Enantiomers (b) Diastereomers (c) Rotamers (d) Homomers

Q.82 For the following compounds A and B the correct statement is



A



B

- (a) A is aromatic and B is antiaromatic. (b) A is antiaromatic and B is non-aromatic.
(c) A and B are both aromatic. (d) A and B are both non-aromatic

Q.83 I for P -31 is:

- (a) 1 (b) $1/2$ (c) $3/2$ (d) 3

Q.84 What is the right order of coupling constants?

- (a) $J^1 > J^2 > J^3$ (b) $J^1 < J^2 < J^3$ (c) $J^1 = J^2 = J^3$ (d) None of these

Q.85 Which aromatic band shows fine structure?

- (a) Primary (b) Secondary (c) Tertiary (d) None

Q.86 Which is a better Diels Alder Diene for reaction with maleic anhydride?

- (a) Furan (b) Pyrrole (c) Thiophene (d) Pyridine

Q.87 Which is a strong base?

- (a) Aniline (b) Cyclohexane (c) Pyrrole (d) Quinoline

Q.88 Which is right order of nucleophilicity?

- (a) $CH_3 - CH_2^\ominus > NH_2^\ominus > CH \equiv C^\ominus > HO^\ominus$ (b) $CH \equiv C^\ominus > NH_2^\ominus > CH_3 - CH_2^\ominus > HO^\ominus$
(c) $HO^\ominus > NH_2^\ominus > CH \equiv C^\ominus > CH_3 - CH_2^\ominus$ (d) $NH_2^\ominus > CH \equiv C^\ominus > HO^\ominus > CH_3 - CH_2^\ominus$

Q.89 Which gives single mono-nitro derivative?

- (a) Naphthalene (b) O-xylene (c) Ethylbenzene (d) P-xylene

Q.90 Which one is most effective in an SN^2 displacement on methyl bromide?

- (a) $C_2H_5O^\ominus$ (b) HO^\ominus (c) $C_6H_5O^\ominus$ (d) CH_3COO^\ominus

Q.91 Which react fastest with N-bromosuccinimide (NBS)?

- (a) Toluene (b) Methane (c) Pyridine (d) Benzene

Q.92 When vinyl cyanide reacts with ethylalcohols in presence of a base, what is formed?

- (a) $CH_2=CH-OH$ (b) $C_2H_5O-CH_2-CH_2CN$ (c) CH_3CH_2OH (d) $C_2H_5-O-C_2H_5$

Q.93 Which is the best leaving group?

- (a) Chloride (b) Fluoride (c) Tosylate (d) None

Q.94 With cis-alkene, the triplet carbene give:

- (a) cis-product (b) trans-product (c) No product (d) Both cis and trans products

Q.95 DNFB is used to identify N-terminal amino acid of peptides. The reagent is called:

- (a) Van-Slyke reagent (b) Sorenson reagent
(c) Sanger's reagent (d) Stephens reagent

Q.96 Continuous wave NMR spectroscopy involves:

- (a) Sequential detection of resonances of nuclei
(b) Simultaneous detection of all resonances of nuclei
(c) Sometimes simultaneous and sometimes sequential detection of resonances of nuclei
(d) None

Q.97 The addition of Br_2 to methyl acetylene to give trans-1,2-dibromopropene is a:

- (a) Stereoselective reaction (b) Stereospecific reaction
(c) Stereoselective and Stereospecific reaction (d) None

Q.98 The reagent used in Edman degradation for N-terminal group analysis of peptides is:

- (a) Phenyl isothiocyanate (b) Benzylchloroformate
(c) DNFB (d) Di-t-butyl carbonate

Q.99 Aspartic acid shows:

- (a) pK_{a1} (b) pK_{a2} (c) pK_{a1} and pK_{a2} (d) pK_{a1} , pK_{a2} and pK_{a3}

Q.100 Which is incorrect about grading of sugars?

- (a) Sucrose-1 (b) Fructose-1.75 (c) Lactose-6 (d) Saccharin-3500

LEGAL NOTICE

This document is an excerpt from the book entitled “MDU M.SC Entrance Chemistry Solved Papers”, and is the intellectual property of the Publisher. The content of this document is protected by international copyright law and is valid only for the personal preview of the user who has originally downloaded it from the publisher’s website (www.dalalinstitute.com). Any act of copying (including plagiarizing its language) or sharing this document will result in severe civil and criminal prosecution to the maximum extent possible under law.



This is a low resolution version only for preview purpose. If you want to read the full book, please consider buying.

Buy the complete book with TOC navigation, high resolution images and no watermark.

Home

CLASSES

NET-JRF, IIT-GATE, M.Sc Entrance & IIT-JAM

Want to study chemistry for CSIR UGC – NET JRF, IIT-GATE, M.Sc Entrance, IIT-JAM, UPSC, ISRO, IISc, TIFR, DRDO, BARC, JEST, GRE, Ph.D Entrance or any other competitive examination where chemistry is a paper ?

[READ MORE](#)

BOOKS

Publications

Are you interested in books (Print and Ebook) published by Dalal Institute ?

[READ MORE](#)

VIDEOS

Video Lectures

Want video lectures in chemistry for CSIR UGC – NET JRF, IIT-GATE, M.Sc Entrance, IIT-JAM, UPSC, ISRO, IISc, TIFR, DRDO, BARC, JEST, GRE, Ph.D Entrance or any other competitive examination where chemistry is a paper ?

[READ MORE](#)

[Home](https://www.dalalinstitute.com/): <https://www.dalalinstitute.com/>

[Classes](https://www.dalalinstitute.com/classes/): <https://www.dalalinstitute.com/classes/>

[Books](https://www.dalalinstitute.com/books/): <https://www.dalalinstitute.com/books/>

[Videos](https://www.dalalinstitute.com/videos/): <https://www.dalalinstitute.com/videos/>

[Location](https://www.dalalinstitute.com/location/): <https://www.dalalinstitute.com/location/>

[Contact Us](https://www.dalalinstitute.com/contact-us/): <https://www.dalalinstitute.com/contact-us/>

[About Us](https://www.dalalinstitute.com/about-us/): <https://www.dalalinstitute.com/about-us/>

Postgraduate Level Classes (NET-JRF & IIT-GATE)

Admission

[Regular Program](#)
[Test Series](#)

[Distance Learning](#)
[Result](#)

Undergraduate Level Classes (M.Sc Entrance & IIT-JAM)

Admission

[Regular Program](#)
[Test Series](#)

[Distance Learning](#)
[Result](#)

MDU M.Sc Entrance Chemistry Solved Papers

“MDU M.Sc Entrance Chemistry Solved Papers” is now available, visit our website for more info.

[READ MORE](#)

Join the revolution by becoming a part of our community and get all of the member benefits like downloading any PDF document for your personal preview.


[Sign Up](#)

Dalal Institute's

MDU M.SC ENTRANCE

Chemistry

Solved Papers



2011-2019 & Letest Model Test




Table of Contents

MDU M.Sc Entrance: Model Test	5
Chemistry	5
❖ Question Paper	5
❖ Answer Key	22
❖ Solution	23
MDU M.Sc Entrance: 2011.....	27
Chemistry	27
❖ Question Paper	27
❖ Answer Key	42
❖ Solution	43
MDU M.Sc Entrance: 2012.....	47
Chemistry	47
❖ Question Paper	47
❖ Answer Key	65
❖ Solution	66
MDU M.Sc Entrance: 2013.....	70
Chemistry	70
❖ Question Paper	70
❖ Answer Key	83
❖ Solution	84
MDU M.Sc Entrance: 2014.....	88
Chemistry	88
❖ Question Paper	88
❖ Answer Key	104
❖ Solution	105
MDU M.Sc Entrance: 2015.....	109
Chemistry	109
❖ Question Paper	109

❖ Answer Key	125
❖ Solution	126
MDU M.Sc Entrance: 2016.....	130
Chemistry	130
❖ Question Paper	130
❖ Answer Key	148
❖ Solution	149
MDU M.Sc Entrance: 2017.....	153
Chemistry	153
❖ Question Paper	153
❖ Answer key	170
❖ Solution	171
MDU M.Sc Entrance: 2018.....	175
Chemistry	175
❖ Question Paper	175
❖ Answer Key	192
❖ Solution	193
MDU M.Sc Entrance: 2019.....	197
Chemistry	197
❖ Question Paper	197
❖ Answer Key	213
❖ Solution	214

The best institute for CSIR-JRF, UGC-NET, IIT-GATE, IIT-JAM, UPSC, GRE, IISc, TIFR, DRDO, BARC, JEST, ISRO and all Ph.D-M.Sc entrance examinations where chemistry is a paper.

***Dalal Institute's
other publications in this series***



**D DALAL
INSTITUTE**

**Main Market, Sector 14, Rohtak, Haryana 124001, India
(info@dalalinstitute.com, +91-9802825820)
www.dalalinstitute.com**